

香港考試局  
HONG KONG EXAMINATIONS AUTHORITY

一九九九年香港中學會考  
HONG KONG CERTIFICATE OF EDUCATION EXAMINATION, 1999

附加數學 試卷二  
ADDITIONAL MATHEMATICS PAPER 2

本評卷參考乃考試局專為今年本科考試而編寫，供閱卷員參考之用。閱卷員在完成閱卷工作後，若將本評卷參考提供其任教會考班的本科同事參閱，本局不表反對，但須切記，在任何情況下均不得容許本評卷參考落入學生手中。學生若索閱或求取此等文件，閱卷員/教師應嚴詞拒絕，因學生極可能將評卷參考視為標準答案，以致但知硬背死記，活剝生吞。這種落伍的學習態度，既不符現代教育原則，亦有違考試着重理解能力與運用技巧之旨。因此，本局籲請各閱卷員/教師通力合作，堅守上述原則。

This marking scheme has been prepared by the Hong Kong Examinations Authority for markers' reference. The Examinations Authority has no objection to markers sharing it, after the completion of marking, with colleagues who are teaching the subject. However, under no circumstances should it be given to students because they are likely to regard it as a set of model answers. Markers/teachers should therefore firmly resist students' requests for access to this document. Our examinations emphasise the testing of understanding, the practical application of knowledge and the use of processing skills. Hence the use of model answers, or anything else which encourages rote memorisation, should be considered outmoded and pedagogically unsound. The Examinations Authority is counting on the co-operation of markers/teachers in this regard.

考試結束後，各科評卷參考將存放於教師中心，供教師參閱。

After the examinations, marking schemes will be available for reference at the Teachers' Centres.



# 只限教師參閱 FOR TEACHERS' USE ONLY

## GENERAL INSTRUCTIONS TO MARKERS

1. It is very important that all markers should adhere as closely as possible to the marking scheme. In many cases, however, candidates would use alternative methods not specified in the marking scheme. Markers should be patient in marking these alternative solutions. In general, a correct alternative solution merits all the marks allocated to that part, unless a particular method is specified in the question.
2. In the marking scheme, marks are classified as follows :
  - 'M' marks – awarded for knowing a correct method of solution and attempting to apply it;
  - 'A' marks – awarded for the accuracy of the answer;
  - Marks without 'M' or 'A' – awarded for correctly completing a proof or arriving at an answer given in the question.
3. In marking candidates' work, the benefit of doubt should be given in the candidates' favour.
4. The symbol (pp-1) should be used to denote marks deducted for poor presentation (p.p.). Note the following points:
  - (a) At most deduct 1 mark for p.p. in each question, up to a maximum of 3 marks for the whole paper.
  - (b) For similar p.p., deduct only 1 mark for the first time that it occurs, i.e. do not penalise candidates twice in the whole paper for the same p.p.
  - (c) In any case, do not deduct any marks for p.p. in those steps where candidates could not score any marks.
  - (d) Note : if the final answers are not expressed in the simplest form, deduct 1 mark for p.p.
  - (e) Some cases in which marks should be deducted for p.p. are specified in the marking scheme. However, the lists are by no means exhaustive. Markers should exercise their professional judgement to give p.p.s in other situations.
5. The symbol (u-1) should be used to denote marks deducted for wrong/no units in the final answers (if applicable). Note the following points:
  - (a) At most deduct 1 mark for wrong/no units for the whole paper.
  - (b) Do not deduct any marks for wrong/no units in case candidate's answer was already wrong.
6. Marks entered in the Page Total Box should be the net total score on that page.
7. In the Marking Scheme, steps which can be omitted are enclosed by dotted rectangles [.....], whereas alternative answers are enclosed by solid rectangles [.....].
8. Unless otherwise specified in the question, numerical answers not given in exact values should not be accepted.
9. Unless otherwise specified in the question, use of notations different from those in the marking scheme should not be penalised.
10. Unless the form of answer is specified in the question, alternative simplified forms of answers different from those in the marking scheme should be accepted if they were correct.

# 只限教師參閱 FOR TEACHERS' USE ONLY

Solution	Marks	Remarks
<p>1.</p> $\int_0^{\frac{\pi}{2}} \cos^2 x dx$ $= \int_0^{\frac{\pi}{2}} \frac{1}{2} (1 + \cos 2x) dx$ $= \left[ \frac{1}{2} \left( x + \frac{1}{2} \sin 2x \right) \right]_0^{\frac{\pi}{2}}$ $= \frac{\pi}{4}$	1A 1A 1A <u>1A</u> <u>3</u>	
<p>2. Let <math>u = x + 2</math>.</p> $\int x(x+2)^{99} dx$ $= \int (u-2) u^{99} du$ $= \int (u^{100} - 2u^{99}) du$ $= \frac{u^{101}}{101} - \frac{u^{100}}{50} + c \quad (c \text{ is a constant})$ $= \frac{(x+2)^{101}}{101} - \frac{(x+2)^{100}}{50} + c$	1M 1A 1A 1A 1A	Accept other suitable substitutions Omit $du$ in most cases (pp-1) Awarded even if $c$ was omitted Withhold this mark if $c$ was omitted
<p><u>Alternative solution</u></p> $\int x(x+2)^{99} dx$ $= \int x \sum_{i=0}^{99} {}_{99}C_i (2^{99-i}) (x^i) dx$ $= \sum_{i=0}^{99} {}_{99}C_i (2^{99-i}) \int x^{i+1} dx$ $= \sum_{i=0}^{99} \frac{{}_{99}C_i (2^{99-i}) x^{i+2}}{(i+2)} + c$ <div style="border: 1px solid black; padding: 5px;"> <math display="block">(OR) = \frac{x^{101}}{101} + {}_{99}C_1 \frac{x^{100}}{50} + {}_{99}C_2 \frac{4x^{99}}{99} + \dots + 2^{98} x^2 + c</math> </div>	1M 3A 3A	For using binomial expansion Should at least contain first two terms and last term. Deduct 1A for each wrong term, up to zero.

# 只限教師參閱 FOR TEACHERS' USE ONLY

Solution	Marks	Remarks
3. (a) The $y$ -intercept of $L_1$ is $\frac{1}{2}$ .	1A	(pp-1) for $(0, \frac{1}{2})$
(b) Distance between $L_1$ and $L_2$ $= \left  \frac{2(0) + 2(\frac{1}{2}) - 13}{\sqrt{2^2 + 2^2}} \right $ $= 3\sqrt{2}$	1M 1A	Accept omitting absolute sign
<b>Alternative solution (1)</b> Distance $= \left  \frac{-13 - (-1)}{\sqrt{2^2 + 2^2}} \right $ $= 3\sqrt{2}$	1M 1A	For using the formula $d = \left  \frac{c_1 - c_2}{\sqrt{a^2 + b^2}} \right $
<b>Alternative solution (2)</b> $y$ -intercept of $L_2 = \frac{13}{2}$ Distance $= (\frac{13}{2} - \frac{1}{2}) \sin 45^\circ$ $= 3\sqrt{2}$	1M 1A	
(c) Let the equation of $L_3$ be $2x + 2y + c = 0$ .		
$\left  \frac{2(0) + 2(\frac{1}{2}) + c}{\sqrt{2^2 + 2^2}} \right  = 3\sqrt{2}$ $ 1 + c  = 12$ $c = 11$ [or $-13$ (rejected)]	1M	<b>OR</b> $c - (-1) = -1 - (-13)$ $c = 11$
$\therefore$ the equation of $L_3$ is $2x + 2y + 11 = 0$ .	1A	Accept equivalent forms
<b>Alternative solution (1)</b> Let the equation of $L_3$ be $2x + 2y + c = 0$ .		
$\left  \frac{c - (-1)}{\sqrt{2^2 + 2^2}} \right  = 3\sqrt{2}$ $ 1 + c  = 12$ $c = 11$ [or $-13$ (rejected)]	1M	
$\therefore$ the equation of $L_3$ is $2x + 2y + 11 = 0$ .	1A	
<b>Alternative solution (2)</b> $y$ -intercept of $L_1 = \frac{1}{2}$ and $y$ -intercept of $L_2 = \frac{13}{2}$ $\therefore$ $y$ -intercept of $L_3 = \frac{1}{2} - (\frac{13}{2} - \frac{1}{2})$ $= -\frac{11}{2}$ Slope of $L_3$ = slope of $L_1 = -1$ $\therefore$ Equation of $L_3$ is $\frac{y - (-\frac{11}{2})}{x} = -1$ $2x + 2y + 11 = 0$	1M	

**只限教師參閱 FOR TEACHERS' USE ONLY**

Solution	Marks	Remarks
<p>Alternative solution (3)</p> <p>Let <math>(x, y)</math> be a point on <math>L_3</math>.</p> $\frac{ 2x + 2y - 1 }{\sqrt{2^2 + 2^2}} = 3\sqrt{2}$ $ 2x + 2y - 1  = 12$ $2x + 2y - 1 = 12 \quad \text{or} \quad 2x + 2y - 1 = -12$ $2x + 2y + 11 = 0 \quad \text{or} \quad 2x + 2y - 13 = 0 \quad (\text{rejected})$ <p><math>\therefore</math> the equation of <math>L_3</math> is <math>2x + 2y + 11 = 0</math>.</p>	1M 1A	
	5	

# 只限教師參閱 FOR TEACHERS' USE ONLY

Solution	Marks	Remarks
<p>4. <u>Marking criteria</u></p> <ul style="list-style-type: none"> <li>- Area <math>A = \int_a^b y dx</math></li> <li>- A correct expression for the shaded area</li> <li>- One correct primitive function</li> <li>- Answer</li> </ul>	<p>1M 2A 1A 1A</p>	<p>A correct expression for the area under a certain portion of a curve – award 1A</p>
$\begin{aligned} \text{Area} &= \int_0^1 (6x^2 - 3x^2) dx + \int_1^2 (6x - 3x^2) dx \\ &= \int_0^1 3x^2 dx + \int_1^2 (6x - 3x^2) dx \\ &= [x^3]_0^1 + [3x^2 - x^3]_1^2 \\ &= 1 + (12 - 8) - (3 - 1) \\ &= 3 \end{aligned}$	<p>1M+2A 1A 1A</p>	<p>Omit <math>dx</math> in most cases (pp-1)</p>
<p><u>Alternative solution (1)</u></p> $\begin{aligned} \text{Area} &= \int_0^2 (6x - 3x^2) dx - \int_0^1 (6x - 6x^2) dx \\ &= [3x^2 - x^3]_0^2 - [3x^2 - 2x^3]_0^1 \\ &= 4 - 1 \\ &= 3 \end{aligned}$	<p>1M+2A 1A 1A</p>	<p>For primitive function, awarded if one was correct</p>
<p><u>Alternative solution (2)</u></p> $\begin{aligned} \text{Area} &= \int_0^1 6x^2 dx + \int_1^2 6x dx - \int_0^2 3x^2 dx \\ &= [2x^3]_0^1 + [3x^2]_1^2 - [x^3]_0^2 \\ &= 2 + 9 - 8 \\ &= 3 \end{aligned}$	<p>1M+2A 1A 1A</p>	
<p><u>Alternative solution (3)</u></p> $\begin{aligned} \text{Area} &= \int_0^6 \left( \sqrt{\frac{y}{3}} - \sqrt{\frac{y}{6}} \right) dy + \int_6^{12} \left( \sqrt{\frac{y}{3}} - \frac{y}{6} \right) dy \\ &= \left[ \frac{2}{3\sqrt{3}} y^{\frac{3}{2}} - \frac{2}{3\sqrt{6}} y^{\frac{3}{2}} \right]_0^6 + \left[ \frac{2}{3\sqrt{3}} y^{\frac{3}{2}} - \frac{y^2}{12} \right]_6^{12} \\ &= 4\sqrt{2} - 4 + (16 - 12) - (4\sqrt{2} - 3) \\ &= 3 \end{aligned}$	<p>1M+2A 1A 1A</p>	
		5

**只限教師參閱 FOR TEACHERS' USE ONLY**

Solution	Marks	Remarks
5. (a) Put $x = 5, y = 0$ : $-3 + k(5+1) = 0$ $k = \frac{1}{2}$ $\therefore$ the equation of $L_1$ is $y - 3 + \frac{1}{2}(x - y + 1) = 0$ $x + y - 5 = 0$ .	1M 1A	Accept equivalent forms
<b>Alternative solution (1)</b> $\begin{cases} y - 3 = 0 \\ x - y + 1 = 0 \end{cases}$  Solve the equations, $x = 2$ and $y = 3$ . The equation of $L_1$ is $\frac{y-3}{x-2} = \frac{0-3}{5-2}$ $x + y - 5 = 0$	1M 1A	
<b>Alternative solution (2)</b> $y - 3 + k(x - y + 1) = 0$ $kx + (1 - k)y + (k - 3) = 0$ Put $y = 0$ : $x = \frac{3 - k}{k} = 5$ $3 - k = 5k$ $k = \frac{1}{2}$ $\therefore$ the equation of $L_1$ is $x + y - 5 = 0$ .	1M 1A	
(b) The equation of $L_2$ is $y - 3 = 0$ .	2A	
<b>Alternative solution</b> $kx + (1 - k)y + (k - 3) = 0$ Slope $= \frac{-k}{1-k} = 0$ $k = 0$ $\therefore$ the equation of $L_2$ is $y - 3 = 0$ .	1M 1A	
(c) Slope of $L_1 = -1$ Let $\theta$ be the acute angle between $L_1$ and $L_2$ . $\tan \theta =  -1 $ $\theta = 45^\circ$ (OR $\frac{\pi}{4}$ )	1M 1A 6	OR $\tan \theta = \left  \frac{-1-0}{1+(-1)(0)} \right $ Accept omitting absolute sign

# 只限教師參閱 FOR TEACHERS' USE ONLY

Solution	Marks	Remarks
6. (a) $\frac{dy}{dx} \Big _{(2, 0)} = 0$ $3(2)^2 - 2(2) + k = 0$ $k = -8$	1M 1A	
(b) $y = \int (3x^2 - 2x - 8) dx$ $= x^3 - x^2 - 8x + c$ <span style="border: 1px dashed black; padding: 2px;"><math>c</math> is a constant</span>	1M 1M	Withhold this mark if "y =" is omitted Awarded even if $c$ is omitted
Put $x = 2, y = 0 : 0 = 2^3 - 2^2 - 8(2) + c$  $c = 12$	1M 1A	For finding $c$
$\therefore$ the equation of the curve is $y = x^3 - x^2 - 8x + 12.$	<u>6</u>	
7. (a) $(1+2x)^n = 1 + {}_n C_1 (2x) + {}_n C_2 (2x)^2 + {}_n C_3 (2x)^3 + \dots$ $= 1 + 2 {}_n C_1 x + 4 {}_n C_2 x^2 + 8 {}_n C_3 x^3 + \dots$ $= 1 + 2nx + 2n(n-1)x^2 + \frac{4}{3}n(n-1)(n-2)x^3 + \dots$	1A 1A	Omit dots (pp-1) in all cases Deduct 1A for each wrong term, up to zero
(b) $(x - \frac{3}{x})^2 (1+2x)^n$ $= (x^2 - 6 + \frac{9}{x^2})(1 + 2 {}_n C_1 x + 4 {}_n C_2 x^2 + \dots)$ $-6 + 36 {}_n C_2 = 210$ ${}_n C_2 = 6$ $\frac{n(n-1)}{2} = 6$ $n^2 - n - 12 = 0$ <span style="border: 1px dashed black; padding: 2px;"><math>(n+3)(n-4) = 0</math></span>	1A 1M 1M	For expanding $(x - \frac{3}{x})^2$
$n = 4$ <span style="border: 1px dashed black; padding: 2px;"><math>\text{or } = -3 \text{ (rejected)}</math></span>	<u>1A</u> <u>6</u>	For ${}_n C_2 = \frac{n(n-1)}{2}$ (OR $= \frac{n!}{2!(n-2)!}$ ) (can be awarded in (a))

# 只限教師參閱 FOR TEACHERS' USE ONLY

Solution	Marks	Remarks
<p>8. (a) <math>\cos 3\theta = \cos(\theta + 2\theta)</math>  <math>= \cos \theta \cos 2\theta - \sin \theta \sin 2\theta</math>  <math>= \cos \theta (2 \cos^2 \theta - 1) - \sin \theta (2 \sin \theta \cos \theta)</math>  <math>= \cos \theta (2 \cos^2 \theta - 1) - 2 \cos \theta (1 - \cos^2 \theta)</math>  <math>= 4 \cos^3 \theta - 3 \cos \theta</math></p>	1A } 1M 1	For expanding $\cos(a+b)$ For expressing in terms of $\cos \theta$
<p><u>Alternative solution (1)</u>  <math>4 \cos^3 \theta - 3 \cos \theta</math>  <math>= \cos \theta (4 \cos^2 \theta - 3)</math>  <math>= \cos \theta [2(1 + \cos 2\theta) - 3]</math>  <math>= 2 \cos \theta \cos 2\theta - \cos \theta</math>  <math>= \cos 3\theta + \cos \theta - \cos \theta</math>  <math>= \cos 3\theta</math></p>	1M 1A 1	For $\cos^2 \theta = \frac{1}{2}(1 + \cos 2\theta)$
<p><u>Alternative solution (2)</u>  <math>(\cos \theta + i \sin \theta)^3 = \cos 3\theta + i \sin 3\theta</math>  <math>(\cos \theta + i \sin \theta)^3 = \cos^3 \theta + 3 \cos^2 \theta (i \sin \theta) +</math>  <math>3 \cos \theta (i \sin \theta)^2 + (i \sin \theta)^3</math></p> <p>Equate real parts :  <math>\cos 3\theta = \cos^3 \theta - 3 \cos \theta (\sin^2 \theta)</math>  <math>= \cos^3 \theta - 3 \cos \theta (1 - \cos^2 \theta)</math>  <math>= 4 \cos^3 \theta - 3 \cos \theta</math></p>	} 1A 1M 1	
<p>(b) <math>\cos 6x + 4 \cos 2x = 0</math>  <math>4 \cos^3 2x - 3 \cos 2x + 4 \cos 2x = 0</math>  <math>4 \cos^3 2x + \cos 2x = 0</math>  <math>\cos 2x = 0 \quad \text{or} \quad 4 \cos^2 2x + 1 = 0</math>  <math>\cos 2x = 0 \quad \boxed{\text{or } \cos^2 2x = -\frac{1}{4} \text{ (rejected)}}</math></p> <p><math>2x = 2k\pi \pm \frac{\pi}{2} \quad \boxed{(k \text{ is an integer})}</math></p> <p><math>x = k\pi \pm \frac{\pi}{4} \quad [\text{OR } x = \frac{1}{4}(2n+1)\pi \quad (n \text{ is an integer})]</math></p>	1A 1A 1M 1A	For $2x = 2n\pi \pm \alpha$ Accept degrees $k\pi \pm 45^\circ$ etc. (pp-1)
<p><u>Alternative solution</u>  <math>\cos 6x + 4 \cos 2x = 0</math>  <math>\cos 6x + \cos 2x + 3 \cos 2x = 0</math>  <math>2 \cos 4x \cos 2x + 3 \cos 2x = 0</math>  <math>\cos 2x = 0 \quad \text{or} \quad 2 \cos 4x + 3 = 0</math>  <math>\cos 2x = 0 \quad \boxed{\text{or } \cos 4x = -\frac{3}{2} \text{ (rejected)}}</math></p> <p><math>2x = 2k\pi \pm \frac{\pi}{2}</math></p> <p><math>x = k\pi \pm \frac{\pi}{4} \quad [\text{OR } x = \frac{1}{4}(2n+1)\pi]</math></p>	1A 1A 1M 1A	For $2x = 2n\pi \pm \alpha$

# 只限教師參閱 FOR TEACHERS' USE ONLY

Solution	Marks	Remarks
9. (a) The equation of $L$ is $y = mx + 1$ . Substitute $y = mx + 1$ into $x^2 = 4y$ : $x^2 = 4(mx + 1)$ $x^2 - 4mx - 4 = 0$ $\therefore x_1, x_2$ are the roots of the equation $x^2 - 4mx - 4 = 0$ .	1A 1M <u>1</u> <u>3</u>	
(b) $\begin{cases} x_1 + x_2 = 4m \\ x_1 x_2 = -4 \end{cases}$ $(x_1 - x_2)^2 = (x_1 + x_2)^2 - 4x_1 x_2$ $= (4m)^2 - 4(-4)$ $= 16(m^2 + 1)$ $AB^2 = (x_1 - x_2)^2 + (y_1 - y_2)^2$ $= (x_1 - x_2)^2 + (mx_1 + 1 - mx_2 - 1)^2$ $= (x_1 - x_2)^2 + (mx_1 - mx_2)^2$ $= (1 + m^2)(x_1 - x_2)^2$ $= (1 + m^2)[16(m^2 + 1)]$ $AB = 4(1 + m^2)$	1A 1M 1A 1M <u>1</u> <u>6</u>	For expressing $y_1, y_2$ in terms of $x_1, x_2$ $\text{OR } (x_1 - x_2)^2 + (\frac{1}{4}x_1^2 - \frac{1}{4}x_2^2)^2$ For using the above result
(c) (i) $x$ -coordinate of centre of $C = \frac{x_1 + x_2}{2}$ $= 2m$ $y$ -coordinate of centre of $C = \frac{y_1 + y_2}{2}$ $= \frac{mx_1 + 1 + mx_2 + 1}{2}$ $= \frac{m}{2}(4m) + 1$ $= 2m^2 + 1$ $\therefore$ the coordinates of the centre are $(2m, 2m^2 + 1)$ . Radius of $C = \frac{AB}{2}$ $= 2(1 + m^2)$	1A 1A 1A 1A	$\boxed{\text{OR } y = m(2m) + 1 = 2m^2 + 1}$
(ii) Distance from centre of $C$ to $y + 1 = 0$ $=  2m^2 + 1 - (-1) $ $\boxed{\text{OR } = \sqrt{(2m^2 + 1) + 1}}$ $= 2(m^2 + 1)$  As the distance from centre of $C$ to $y + 1 = 0$ is equal to the radius of $C$ , the line $y + 1 = 0$ is a tangent to $C$ . $\boxed{\text{OR The line } y + 1 = 0 \text{ and } C \text{ meet at one point.}}$	1M 1A  $\left\{ \begin{array}{l} 1M+1A \\ \hline 7 \end{array} \right.$	Accept omitting absolute sign 1M for comparing the radius and the distance

# 只限教師參閱 FOR TEACHERS' USE ONLY

Solution	Marks	Remarks
10. (a) $PA = \sqrt{3}PB$ $\sqrt{(x+3)^2 + y^2} = \sqrt{3} \sqrt{(x+1)^2 + y^2}$ $x^2 + 6x + 9 + y^2 = 3(x^2 + 2x + 1 + y^2)$ $x^2 + y^2 = 3$	1A 1M <hr/> 1 <hr/> 3	(can be omitted) For squaring and expanding both sides
(b) Differentiate $x^2 + y^2 = 3$ with respect to $x$ : $2x + 2y \frac{dy}{dx} = 0$ $\frac{dy}{dx} = -\frac{x}{y}$ Equation of tangent at $T(a, b)$ is $\frac{y-b}{x-a} = \frac{-a}{b}$ $by - b^2 = -ax + a^2$ $ax + by = a^2 + b^2$	<hr/> 1M	
<div style="border: 1px solid black; padding: 2px; display: inline-block;"> <math>\text{QR } ax + by = 3</math> </div>	1A	
<b>Alternative solution</b> Using the formula $xx_1 + yy_1 = 3$ , the equation of the tangent at $T$ is $ax + by = 3$ .	2A	<hr/> 2
(c) Substitute $A(-3, 0)$ into the equation of tangent: $a(-3) + b(0) = 3$ $a = -1$ $b = \sqrt{3 - (-1)^2}$ <span style="border: 1px dashed black; padding: 2px;">(<math>\because S</math> lies in the 2nd quadrant.)</span> $= \sqrt{2}$ $\therefore$ the coordinates of $S$ are $(-1, \sqrt{2})$ .	1M 1A 1A	
<b>Alternative solution</b> 		
Let $\phi$ be the angle between $OS$ and the negative $x$ -axis. $\angle OSA = \frac{\pi}{2}$ $-x = OS \cos \phi$ $= \sqrt{3} \left(\frac{\sqrt{3}}{3}\right)$ $x = -1$ $y = OS \sin \phi$ $= \sqrt{3} \left(\frac{\sqrt{6}}{3}\right)$ $= \sqrt{2}$ $\therefore$ the coordinates of $S$ are $(-1, \sqrt{2})$ .	1M 1A 1A	<hr/> 3


# 只限教師參閱 FOR TEACHERS' USE ONLY

Solution	Marks	Remarks						
(d) (i) The coordinates of $Q$ are $(-3+r \cos \theta, r \sin \theta)$ .	1A+1A							
(ii) (1) Substitute $(-3+r \cos \theta, r \sin \theta)$ into $C$ :	1M							
$(-3+r \cos \theta)^2 + (r \sin \theta)^2 = 3$								
$9 - 6r \cos \theta + r^2 \cos^2 \theta + r^2 \sin^2 \theta = 3$								
$r^2 - 6r \cos \theta + 6 = 0 \quad \dots (*)$								
Since $AH = r_1$ , $AK = r_2$ , $r_1$ and $r_2$ are the roots of $(*)$ .	1							
(2) Since $\ell$ cuts $C$ at two distinct points, $(*)$ has two distinct real roots.								
$(6 \cos \theta)^2 - 4(6) > 0$	1M							
$\cos^2 \theta > \frac{2}{3}$	1A							
$\cos \theta > \sqrt{\frac{2}{3}} \quad \text{or} \quad \cos \theta < -\sqrt{\frac{2}{3}}$	1A							
<div style="border: 1px dashed black; padding: 5px;"><math>\left(\text{rejected } \because -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}\right)</math></div>		Can be awarded even if considering $\Delta = 0$ or $\Delta \geq 0$						
$\therefore -0.615 < \theta < 0.615$ (correct to 3 sig. figures)	1A	(OR $-35.3^\circ < \theta < 35.3^\circ$ )						
<b>Alternative solution</b> Let $\alpha$ be the angle between $AS$ and the $x$ -axis.								
$\tan \alpha = \frac{SB}{AB}$ <table border="1" style="display: inline-table; vertical-align: middle;"> <tr> <td style="padding: 5px;">OR</td> <td style="padding: 5px;"><math>\sin \alpha = \frac{OS}{OA}</math></td> <td style="padding: 5px;"><math>\sin \alpha = \frac{SB}{SA}</math></td> </tr> <tr> <td style="padding: 5px;"><math>= \frac{\sqrt{2}}{2}</math></td> <td style="padding: 5px;"><math>= \frac{\sqrt{3}}{3}</math></td> <td style="padding: 5px;"><math>= \frac{\sqrt{2}}{\sqrt{6}} = \frac{1}{\sqrt{3}}</math></td> </tr> </table>	OR	$\sin \alpha = \frac{OS}{OA}$	$\sin \alpha = \frac{SB}{SA}$	$= \frac{\sqrt{2}}{2}$	$= \frac{\sqrt{3}}{3}$	$= \frac{\sqrt{2}}{\sqrt{6}} = \frac{1}{\sqrt{3}}$	1M	
OR	$\sin \alpha = \frac{OS}{OA}$	$\sin \alpha = \frac{SB}{SA}$						
$= \frac{\sqrt{2}}{2}$	$= \frac{\sqrt{3}}{3}$	$= \frac{\sqrt{2}}{\sqrt{6}} = \frac{1}{\sqrt{3}}$						
$\alpha = 35.3^\circ$ (correct to 3 sig. figures)	1A							
Since $\ell$ cuts the circle at two distinct points, $-0.615 < \theta < 0.615$ .	1A							
		<b>OR</b> $S$ is a point on the locus $\therefore SA = \sqrt{3} SB$ $\therefore \sin \alpha = \frac{SB}{SA} = \frac{1}{\sqrt{3}}$						

# 只限教師參閱 FOR TEACHERS' USE ONLY

Solution	Marks	Remarks
11. (a) Consider $\triangle ABD$ :		
By Sine Law,		
$\frac{AD}{\sin \angle ABD} = \frac{\ell}{\sin \angle ADB}$	1M	
$\frac{AD}{\sin(180^\circ - \alpha)} = \frac{\ell}{\sin(\alpha - 10^\circ)}$		
$AD = \frac{\ell \sin \alpha}{\sin(\alpha - 10^\circ)} \text{ m}$	1	
(i) Consider $\triangle ACD$ :		
$CD = AD \sin 10^\circ$		
$= \frac{\ell \sin \alpha \sin 10^\circ}{\sin(\alpha - 10^\circ)} \text{ m}$	1A	
(ii) Consider $\triangle ADH$ :		
$\frac{AD}{\sin(\alpha - \beta)} = \frac{DH}{\sin(\beta - 10^\circ)}$	1M	
$DH = AD \frac{\sin(\beta - 10^\circ)}{\sin(\alpha - \beta)} = \frac{\ell \sin \alpha \sin(\beta - 10^\circ)}{\sin(\alpha - 10^\circ) \sin(\alpha - \beta)}$		
Consider $\triangle DHG$ :		
$h = DH \sin \alpha$		
$= \frac{\ell \sin^2 \alpha \sin(\beta - 10^\circ)}{\sin(\alpha - 10^\circ) \sin(\alpha - \beta)}$	1	
	6	
(b) (i) (1) Using (a) (ii) :		
height of pole $= \frac{97 \sin^2 15^\circ \sin(10.2^\circ - 10^\circ)}{\sin(15^\circ - 10^\circ) \sin(15^\circ - 10.2^\circ)}$		
$= 3.1100$		
$= 3.1 \text{ m (correct to 2 sig. fig.)}$	1A	Omit/wrong unit ( $u - 1$ )
(2) Using (a) (i) :		
height of tower $CD = \frac{97 \sin 15^\circ \sin 10^\circ}{\sin(15^\circ - 10^\circ)}$		
$= 50.020$		
$= 50 \text{ m (correct to 2 sig. fig.)}$	1A	
radius of tower $= \frac{h}{\tan 15^\circ}$		
$= \frac{3.1100}{\tan 15^\circ}$	1M	
$= 11.607$		
$= 12 \text{ m (correct to 2 sig. fig.)}$	1A	

只限教師參閱 FOR TEACHERS' USE ONLY

Solution	Marks	Remarks
(ii) (1)		
		
Consider $\Delta HPO$ :		
$\tan \angle HPO = \frac{OH}{OP}$	2M	
$OP = \frac{OH}{\tan 15^\circ}$ $= \frac{3.1100 + 50.020}{\tan 15^\circ}$ $= 198.28$ $= 200 \text{ m (correct to 2 sig. fig.)}$	1M 1A	For $OH = h + CD$
<b>Alternative solution</b> $OP = OB$ $OP = OC + CB$ $= r + \frac{CD}{\tan \alpha}$ $= 11.607 + \frac{50.020}{\tan 15^\circ}$ $= 198.28$ $= 200 \text{ m (correct to 2 sig. fig.)}$	1M 1M 1M 1A	(can be omitted)
$(2) \angle BPO = \frac{1}{2}(180^\circ - 45^\circ)$ $= 67.5^\circ$ <p>Bearing of B from P is N(67.5° - 45°)W, i.e. N22.5°W. (OR N23°W correct to 2 sig. fig.)</p>	1A <hr/> 1A <hr/> 10	(Awarded 1M for other correct methods) 337.5° (OR 340°)

# 只限教師參閱 FOR TEACHERS' USE ONLY

Solution	Marks	Remarks
<p>12. (a) For <math>n = 1</math>, LHS = <math>\cos\theta</math>.</p> $\begin{aligned} \text{RHS} &= \frac{\sin 2\theta}{2 \sin \theta} \\ &= \frac{2\sin\theta\cos\theta}{2\sin\theta} = \cos\theta = \text{LHS}. \end{aligned}$ <p><math>\therefore</math> the statement is true for <math>n = 1</math>.</p> <p>Assume <math>\cos\theta + \cos 3\theta + \cos 5\theta + \dots + \cos(2k-1)\theta = \frac{\sin 2k\theta}{2 \sin \theta}</math> [for some +ve integer <math>k</math>.]</p> <p>Then <math>\cos\theta + \cos 3\theta + \cos 5\theta + \dots + \cos(2k-1)\theta + \cos[2(k+1)-1]\theta</math></p> $\begin{aligned} &= \frac{\sin 2k\theta}{2 \sin \theta} + \cos(2k+1)\theta \\ &= \frac{\sin 2k\theta + 2 \sin \theta \cos(2k+1)\theta}{2 \sin \theta} \\ &= \frac{\sin 2k\theta + \sin(2k+2)\theta - \sin 2k\theta}{2 \sin \theta} \\ &= \frac{\sin 2(k+1)\theta}{2 \sin \theta} \end{aligned}$ <p>The statement is also true for <math>n = k+1</math> if it is true for <math>n = k</math>. By the principle of mathematical induction,</p> <p>the statement is true for all positive integers <math>n</math>.</p>	1 1 1 1 1 <hr/> 1 1	(cannot be omitted)
<p>(b) Using (a) : <math>\cos\theta + \cos 3\theta + \cos 5\theta = \frac{\sin 6\theta}{2 \sin \theta}</math>, where <math>\sin \theta \neq 0</math>.</p> <p>Put <math>\theta = \frac{\pi}{2} - x</math> :</p> $\begin{aligned} \cos\left(\frac{\pi}{2} - x\right) + \cos 3\left(\frac{\pi}{2} - x\right) + \cos 5\left(\frac{\pi}{2} - x\right) &= \frac{\sin 6\left(\frac{\pi}{2} - x\right)}{2 \sin\left(\frac{\pi}{2} - x\right)} \\ \cos\left(\frac{\pi}{2} - x\right) + \cos\left(\frac{3\pi}{2} - 3x\right) + \cos\left(\frac{5\pi}{2} - 5x\right) &= \frac{\sin(3\pi - 6x)}{2 \sin\left(\frac{\pi}{2} - x\right)} \end{aligned}$ $\sin x - \sin 3x + \sin 5x = \frac{\sin 6x}{2 \cos x}$ <p>(where <math>\sin\left(\frac{\pi}{2} - x\right) = \cos x \neq 0</math>)</p> <p><u>Alternative solution</u> Consider <math>2 \cos x (\sin x - \sin 3x + \sin 5x)</math></p> $\begin{aligned} &= \sin 2x - (\sin 4x + \sin 2x) + (\sin 6x + \sin 4x) \\ &= \sin 6x \end{aligned}$ <p><math>\therefore \sin x - \sin 3x + \sin 5x = \frac{\sin 6x}{2 \cos x}</math>, where <math>\cos x \neq 0</math>.</p>	1A <hr/> 1 <hr/> 1	(can be omitted)
	<hr/> <hr/> <hr/>	

# 只限教師參閱 FOR TEACHERS' USE ONLY

Solution	Marks	Remarks
<p>(c)</p> $\int_{0.1}^{0.5} \left( \frac{\sin x - \sin 3x + \sin 5x}{\cos x + \cos 3x + \cos 5x} \right)^2 dx$ $= \int_{0.1}^{0.5} \left[ \frac{\sin 6x}{2 \cos x} / \frac{\sin 6x}{2 \sin x} \right]^2 dx$ $= \int_{0.1}^{0.5} \tan^2 x dx$ $= \int_{0.1}^{0.5} (\sec^2 x - 1) dx$ $= [\tan x - x]_{0.1}^{0.5}$ $= 0.046 \text{ (correct to 2 sig. fig.)}$	1A  1M  1A  1A <hr/> 4	For integrand only  For $\tan^2 x = \sec^2 x - 1$  For primitive function only
<p>(d)</p> $\int_{\frac{\pi}{3}}^{\frac{\pi}{2}} (\sin x + 3 \sin 3x + 5 \sin 5x + 7 \sin 7x + \dots + 1999 \sin 1999x) dx$ $= [-\cos x - \cos 3x - \cos 5x - \cos 7x - \dots - \cos 1999x]_{\frac{\pi}{3}}^{\frac{\pi}{2}}$ $= -\frac{1}{2} \left[ \frac{\sin 2000x}{\sin x} \right]_{\frac{\pi}{3}}^{\frac{\pi}{2}}$ $= -\frac{1}{2} \left( \frac{\sin 1000\pi}{\sin \frac{\pi}{2}} - \frac{\sin \frac{2000\pi}{3}}{\sin \frac{\pi}{3}} \right)$ $= \frac{1}{2}$	1M+1A  1A  1A  1A	1M for integrating each term (At least two terms)
<p><u>Alternative solution</u></p> $\int_{\frac{\pi}{3}}^{\frac{\pi}{2}} (\sin x + 3 \sin 3x + 5 \sin 5x + 7 \sin 7x + \dots + 1999 \sin 1999x) dx$ $= [-\cos x - \cos 3x - \cos 5x - \cos 7x - \dots - \cos 1999x]_{\frac{\pi}{3}}^{\frac{\pi}{2}}$ $= [-\cos \frac{\pi}{2} - \cos \frac{3\pi}{2} - \cos \frac{5\pi}{2} - \dots - \cos \frac{1999\pi}{2}] -$ $[-\cos \frac{\pi}{3} - \cos \pi - \cos \frac{5\pi}{3} - \cos \frac{7\pi}{3} - \dots - \cos \frac{1999\pi}{3}]$ $= 0 - [(-\frac{1}{2} + 1 - \frac{1}{2}) + (-\frac{1}{2} + 1 - \frac{1}{2}) + \dots + (-\frac{1}{2} + 1 - \frac{1}{2}) - \dots]$ $= \frac{1}{2}$	1M+1A  1A  1A <hr/> 4	1M for integrating each term  (can be omitted)

**只限教師參閱 FOR TEACHERS' USE ONLY**

Solution	Marks	Remarks
13. (a) Substitute $(r, 2)$ into $x = \sqrt{4 + 3y^2}$ : $\begin{aligned} r &= \sqrt{4 + 3(2)^2} \\ &= 4 \end{aligned}$	1M 1A <hr/> 2	
(b) $V =$ Volume of lower cylindrical part + volume of upper part $\text{Volume of lower cylindrical part} = \pi r^2 h$ $\begin{aligned} &= \pi(4)^2(2) \\ &= 32\pi \end{aligned}$	1M 1A	
<u>Alternative solution</u> $\text{Volume of lower cylindrical part} = \pi \int_0^2 x^2 dy$ $\begin{aligned} &= \pi \int_0^2 4^2 dy \\ &= 32\pi \end{aligned}$	1M 1A	
$\text{Volume of upper part} = \pi \int_2^h x^2 dy$ $\begin{aligned} &= \pi \int_2^h (4+3y^2) dy \\ &= \pi[4y + y^3]_2^h \\ &= (h^3 + 4h - 16)\pi \\ \therefore V &= 32\pi + (h^3 + 4h - 16)\pi \\ &= (h^3 + 4h + 16)\pi \quad \boxed{\text{cubic units}} \end{aligned}$	1M 1A 1A 1A <hr/> 1 <hr/> 7	For $\pi[4y + y^3]$ only
(c) (i) Let $h$ units be the depth of water at time $t$ . $\frac{dV}{dt} = \pi(3h^2 + 4) \frac{dh}{dt}$ Put $\frac{dV}{dt} = -2\pi$ and $h = 3$ : $-2\pi = \pi[3(3)^2 + 4] \frac{dh}{dt}$ $\frac{dh}{dt} = \frac{-2}{31} \text{ units per sec. (OR } s^{-1}\text{)}$ (OR The depth decreases at a rate $\frac{2}{31}$ units per sec.)	1M+1A  1M  1A	1 M for chain rule  For substitution  (Accept substitute $\frac{dV}{dt} = 2\pi$ ) omit/wrong unit (u-1)

**只限教師參閱 FOR TEACHERS' USE ONLY**

Solution	Marks	Remarks
(ii) When $h = 1$ , the water remained is in the cylindrical part only.		
$\frac{dh}{dt} = \frac{\frac{dV}{dt}}{\text{base area of cylinder}}$	1M	(can be omitted)
$= \frac{-2\pi}{\pi(4)^2}$	1M	For substitution (Accept substitute $\frac{dV}{dt} = 2\pi$ )
$= -\frac{1}{8} \text{ units per sec. (OR } s^{-1}\text{)}$  (OR The depth decreases at a rate $\frac{1}{8}$ units per sec.)	1A	
<b>Alternative solution</b> $V = \pi(4)^2 h$ $= 16\pi h$ $\frac{dV}{dt} = 16\pi \frac{dh}{dt}$ $-2\pi = 16\pi \frac{dh}{dt}$ $\therefore \frac{dh}{dt} = -\frac{1}{8} \text{ units per sec. (OR } s^{-1}\text{)}$	1M 1M 1A	
	7	